

Targeting the Ataxia Telangiectasia and Rad3 Signaling Pathway to Overcome Chemoresistance in Cancer

Marwan Kwok^{*,**}, Tatjana Stankovic^{*}

^{*}Institute of Cancer and Genomic Sciences, University of Birmingham, United Kingdom

^{**}Centre for Clinical Haematology, Queen Elizabeth Hospital Birmingham, United Kingdom

Abstract

The ataxia telangiectasia and Rad3-related (ATR) signaling pathway controls DNA replication initiation and cell-cycle progression and coordinates the repair of damaged replication forks. It is, therefore, an important regulator of the cellular response to DNA replication stress. Inhibition of the ATR signaling pathway results in dysregulation of replication origin initiation, leading to exacerbation of replication stress manifested by the slowing and stalling of replication forks, and enhanced susceptibility to fork breakage. Accordingly, inhibition of ATR or its downstream effector Chk1 is potentially cytotoxic to tumor cells with heightened levels of endogenous replication stress, particularly in tumors with DNA damage response or cell-cycle checkpoint defects. This results in a functional addiction to the ATR pathway. Small-molecule inhibitors of ATR and Chk1 are currently under investigation. Preclinical studies of these inhibitors have demonstrated single-agent efficacy across a range of malignancies, and results from clinical trials are emerging. In addition, markers of sensitivity to ATR and Chk1 inhibitors are being identified, and novel chemotherapeutic combinations involving these agents are being developed. A number of unresolved issues notwithstanding, the ATR signaling pathway represent a promising target for cancer therapeutics.

ABBREVIATIONS

ALT	Alternative lengthening of telomeres
AML	Acute myeloid leukemia
ARID1A	AT-rich interaction domain1A
ATM	Ataxia telangiectasia mutated
ATR	Ataxia telangiectasia and Rad3-related
ATRIP	ATR-interacting protein
BLM	Bloom syndrome protein
BRCA1	Breast cancer type 1 susceptibility protein
cdk	Cyclin-dependent kinase
chk1	Checkpoint kinase 1
CLL	Chronic lymphocytic leukemia

Targeting Cell Survival Pathways to Enhance Response to Chemotherapy. <http://dx.doi.org/10.1016/B978-0-12-813753-6.00010-X>

Copyright © 2018 Elsevier Inc. All rights reserved.

DDR	DNA damage response
DLBCL	Diffuse large B cell lymphoma
DSBs	Double-stranded breaks
FANCM	Fanconi anemia complementation group M
GCB	Germinal center B-cell like
HRR	Homologous recombination repair
NHEJ	Nonhomologous end joining
PARP	Poly-(ADP-ribose) polymerase
PDX	Patient-derived xenotransplantation
PIKK	Phosphoinositide 3-kinase-related kinase
RPA	Replication protein A
RS	Replication stress
ssDNA	Single-stranded DNA
T-ALL	T-cell acute lymphoblastic leukemia
TOPB1	Topoisomerase 2-binding protein 1
WRN	Werner syndrome ATP-dependent helicase

10.1 INTRODUCTION

The DNA damage response (DDR) is essential for the maintenance of genome integrity. This is accomplished through a network of intricately connected pathways, resulting in a dichotomous outcome: either resolution of the incipient DNA damage or apoptotic cell death, depending on the nature and severity of the afflicting lesion. Tumorigenesis is associated with a pervasive corruption of one or more of these pathways, often through mutation or deletion of key DNA repair or regulatory proteins, conferring on malignant cells a survival advantage, avoidance of apoptosis, and chemoresistance [1]. However, the subversion of normal DDR mechanisms can render tumor cells susceptible to further DNA insults, the accumulation of which may become incompatible with survival and results in cell death during mitosis, a process termed mitotic catastrophe [2,3]. Moreover, the disruption of a DNA repair pathway constrains tumor cells to rely on collateral repair pathways to maintain genome stability, thereby creating a vulnerability that can be amenable to therapeutic targeting [4].

Ataxia telangiectasia and Rad3-related (ATR) is a phosphoinositide 3-kinase-related kinase (PIKK) that is the master regulator of one such DDR pathway, with wide-ranging roles in DNA replication [5]. Studies in the last decade have shed light into its function and interaction with other DDR pathways, as well as the effects of inhibiting this pathway. We herein review our current understanding of the mechanistic basis of ATR signaling and illustrate how some tumor cells are distinctively reliant on this pathway and how targeting this pathway leads to tumor cell death. We also highlight important preclinical and clinical studies on this therapeutic target, as a single agent and in combination with other chemotherapeutic agents, and discuss unresolved questions and directions for future research.

10.2 FUNCTIONAL ROLES OF THE ATR SIGNALING CASCADE

10.2.1 The Role of ATR in the Regulation of DNA Replication Initiation

Initiation of DNA replication is a stochastic event, occurring at replication origins from which replication forks arise. Replication origins are clustered and replicated together within

replication factories. An important function of ATR is its ability to protect replication forks by acting as a negative regulator of replication origin initiation. This limits the activation of new replication factories, thereby directing replication toward already active factories [6,7]. Such negative regulation of replication initiation is of paramount importance in preventing the depletion of cellular pools of nucleotides and replication proteins, which would otherwise result in genomic instability. Indeed, only a small proportion of licensed origins is active in an unperturbed cell cycle, with the rest remaining dormant [6].

10.2.2 The Role of the ATR Signaling Pathway in the Maintenance of Replication Fork Stability

During DNA replication when the double-stranded DNA helix unwinds, replication fork progression can slow or stall due to replication obstacles such as DNA damage. Continued unwinding of the DNA helix despite stalled DNA polymerase leads to exposure of long stretches of single-stranded DNA (ssDNA) that are prone to damage. This results in replication fork breakage. The ATR pathway functions to prevent this outcome by protecting exposed ssDNA and repairing DNA lesions at these sites. Exposed ssDNA is rapidly coated with replication protein A (RPA), which stabilizes the replication fork and protects ssDNA from deleterious degradation by nucleases [8].

The ATR signaling cascade is initiated by the independent recruitment of two complexes, namely the ATR and ATR-interacting protein (ATRIP) and the RAD9–RAD1–HUS1 (9–1–1) complex, to RPA-coated ssDNA where RPA binds ATRIP and engages with the 9–1–1 complex. Further recruitment of topoisomerase 2-binding protein 1 (TOPBP1) activates ATR by stimulating ATR kinase activity [9]. Recently, the RPA-binding protein Ewing's tumor-associated antigen 1 was also reported to have ATR stimulatory activity and ability to activate ATR independent of TOPBP1 in a parallel pathway [10–12]. Functionally active ATR in turn phosphorylates the downstream effector molecule Chk1 and other substrates such as Werner syndrome ATP-dependent helicase (WRN), Bloom syndrome protein (BLM), and SMAR-CAL1. A recent study using iPOND method confirmed the recruitment of these proteins to DNA replisomes upon replication fork stalling and highlighted their functional importance in preventing fork collapse and breakage during DNA replication [13].

Through these proteins, ATR promotes replication fork stability and mediates replication restart in the event of fork stalling. This is achieved through fork reversal, whereby stalled forks reverse their course and through fork remodeling resulting in the formation of four-way junctions [14,15]. Helicases such as WRN and BLM, as well as translocases such as SMAR-CAL1 and the Fanconi anemia complementation group M protein (FANCM) are thought to play a major role in this process [16–21]. Fork reversal, remodeling, and restart together ensure completion of DNA replication and prevent fork breakage.

Proteins downstream of ATR also serve additional functions in facilitating fork repair, through suppression of new origin firing (BLM and FANCM) and protection of replication forks from unscheduled DNA degradation (WRN) [22,23]. Furthermore, Chk1 diffuses throughout the nucleus to suppress new origin firing. Thus, the regulatory activities of ATR signaling are amplified and propagated globally in the event of replication fork stalling [6,24]. This is crucial in curtailing further fork stalls to conserve the supply of RPA for the hitherto perturbed sites and to ensure that an excess of RPA over ssDNA is maintained [25].

10.2.3 The Role of the ATR Signaling Pathway in Cell-Cycle Regulation

Finally, through the downstream kinase Chk1, ATR also plays a major role in controlling cell-cycle progression. Chk1 regulates cellular entry into mitosis through the G2/M checkpoint. Mechanistically, this involves phosphorylation of Cdc25 phosphatases to prevent its binding to the cyclin B1–cyclin-dependent kinase (Cdk1) complex. Cdc25 phosphatases remove inhibitory phosphorylation from Cdk1 that is a prerequisite for the activation of cyclin B1–Cdk1, which initiates mitotic entry. In the presence of stalled replication forks, abolition of cyclin B1–Cdk1 activity through Chk1-mediated phosphorylation of Cdc25 phosphatases prevents premature entry of replicatively perturbed cells into mitosis until replication stalling is rectified and genomic duplication complete [26,27].

10.3 THE ATR PATHWAY AS A CANCER THERAPEUTIC TARGET

10.3.1 Induction of Replication Stress as a Consequence of ATR Pathway Inhibition

When replication fork progression is disrupted, slowing or stalling of replication forks ensues. This results in replication stress (RS), with cells continuing to progress through cell cycle despite the presence of unreplicated DNA [28]. Low levels of RS are ubiquitous in cycling cells and are often tolerated [29]. On the other hand, excessive amounts of RS arising, for instance, from the accumulation of unrepaired DNA damage, misincorporated ribonucleotides, or from difficult-to-replicate segments of DNA within fragile sites, inevitably lead to cell death [28]. The ATR signaling pathway ameliorates RS through its role in the regulation of DNA replication initiation, replication fork stability, and cell-cycle progression, as outlined above. Conversely, inhibition of the ATR pathway is a potent inducer of RS.

A consequence of ATR pathway inhibition is excessive replication origin initiation in an unscheduled and uncontrolled manner. This occurs independent of any additional sources of RS, either endogenous or exogenous. Indeed, DNA fiber analysis showed that the number of active replication forks in cancer cells was increased when ATR is inhibited, even in the absence of DNA damaging agents or other sources of RS [18,30]. Excessive and uncontrolled origin initiation as a result of ATR inhibition depletes essential replication factors such as nucleotides and replication proteins. This leads to a decrease in the rate of replication fork elongation and stalling of replication forks [24] and, as evidenced in cancer cells, to rapid reduction in inter-origin distance indicating increased replication origins [18,31]. These observations are consistent with the induction of RS.

Excessive and uncontrolled initiation of dormant replication origins also generates increased ssDNA. Exposed ssDNA is normally coated with RPA. However, when ATR is inhibited, the excess of ssDNA exhausts the nuclear pool of RPA. Unprotected ssDNA is susceptible to fork breakage, thereby precipitating DNA damage [30]. Furthermore, ATR inhibition prevents proper regulation of SMARCA1, the deregulated activities of which promote excessive replication fork regression and fork collapse [18]. Together, these mechanisms underpin the cytotoxic effect of ATR pathway inhibition.

10.3.2 Replication Stress as a Therapeutic Vulnerability in Cancer Cells

An elevated level of RS is a defining feature of malignant cells. Although the precise mechanisms underlying oncogene-induced RS remain unclear, it is plausible that the accelerated consumption of RPA resulting from unchecked proliferation mediated by oncogenes could be a contributing factor [30]. Other potential mechanisms, such as the deregulation of replication initiation as well as interference between replication and transcription, arising either directly or indirectly from oncogenic activity, have also been postulated [32,33]. Given that the high constitutive levels of cellular RS is inherent in many tumors, it follows that they are likely to be particularly sensitive to RS overload instigated by inhibition of the ATR signaling pathway.

Until recently, it has been assumed that ATR is physiologically indispensable and not amenable to therapeutic targeting [34]. Indeed, human ATR mutations are uncommon, even in cancer. Among the few cancer types where *ATR* mutations have been reported, the acquisition of these mutations has been associated with tumor progression and poor clinical outcome [35,36]. Moreover, abrogation of the *ATR* gene is embryonically lethal, whereas its deletion in adult mice results in rapid aging and stem-cell loss [37,38]. In patients with Seckel syndrome in whom ATR signaling is defective due to hypomorphic germline mutation of the *ATR* gene, growth retardation, dwarfism, microcephaly, and mental impairment are typical manifestations [39,40].

Nonetheless, although complete abolition of ATR activity likely results in toxicity to healthy tissues, suppression of ATR activity was shown to be tolerable in healthy cells but not in tumor cells. In an elegant study by Schoppy et al. [41], conditional reduction of ATR in mice to 10% of normal levels using Cre-lox recombination resulted in minimal adverse effect on healthy hematopoietic and intestinal tissues. In contrast, suppression of ATR to this level was sufficient to severely restrict the growth of fibrosarcomas driven by H-Ras and p53 loss, as well as acute myeloid anemia driven by K-Ras and the *MLL-ENL* translocation [41]. Such reduction in ATR levels likewise prevented the development of Myc-driven lymphomas and pancreatic tumors featuring high levels of RS, as demonstrated by Murga et al. [42]. Corroborating these reports, we recently showed that therapeutic doses of an ATR kinase inhibitor were well tolerated in mice, while achieving effective tumor reduction [31]. Therefore, healthy and tumor cells demonstrate differential sensitivity to ATR pathway inhibition, providing a therapeutic window for the targeting of tumor cells.

10.4 PREDICTIVE BIOMARKERS OF SENSITIVITY TO ATR PATHWAY INHIBITION

10.4.1 ATR Pathway Addiction and the Concept of Synthetic Lethality

The substantial functional redundancy within cellular DDR pathways is widely recognized, wherein more than one pathway is capable of performing the same role. Physiologically, this provides protection against disruption of normal DDR mechanisms, particularly in cancer where DDR genes are mutated with high frequency. However, when one pathway is disrupted, cells become greatly reliant on collateral pathways. Thus, where two independent

pathways regulate an essential DDR process, the absence of one pathway is compatible with cell survival, whereas the absence of both results in cell death. This gives rise to an emerging therapeutic strategy known as synthetic lethality, in which collaborating pathways are abolished to induce cytotoxicity [43,44].

From the studies highlighted above, it is apparent that tumors with heightened levels of RS, such as those driven by Myc or Ras, are exquisitely sensitive to ATR pathway inhibition. Understanding other circumstances under which tumor cells become addicted to the ATR signaling pathway is therefore essential to identifying further predictive biomarkers of sensitivity to ATR inhibition. In the following sections, we summarize current evidence for synthetic lethality of ATR pathway inhibition with various DDR defects prevalent in cancers.

10.4.2 Synthetic Lethality With G1/S Cell-Cycle Checkpoint Defects

The G1/S cell-cycle checkpoint is regulated by p53, a key regulatory transcription factor. In response to DNA damage, p53 transcriptionally activates certain genes and represses others to promote G1/S cell-cycle arrest [45]. The *TP53* tumor suppressor gene encoding p53 is among the most frequently mutated in cancer [46]. The loss of the G1/S checkpoint due to *TP53* loss imposes upon tumor cells a dependence on the G2/M checkpoint controlled primarily through ATR/Chk1. Consequently, ATR pathway inhibition has synthetically lethal properties in *TP53*-defective tumors, the evidence of which can be gleaned from several studies.

Nghiem et al. [47] provided one of the earliest evidence for this concept demonstrating hypersensitivity of G1/S checkpoint-deficient cells to ATR loss. In this study, the authors observed that the osteosarcoma cell line U2OS in the absence of ATR function displayed premature chromatin condensation, a feature associated with mitotic catastrophe, in response to hydroxyurea or ultraviolet radiation-induced RS. Moreover, repression of p53 function by overexpression of MDM2 markedly potentiated the lethal effect of ATR inhibition in these cells [47]. These findings were recapitulated in a later study showing that a p53-deficient colorectal cell line rendered ATR deficient by knock-in of a Seckel gene (*ATR^{se}*) exhibited marked sensitivity to hydroxyurea- and cisplatin-induced RS, whereas restoration of p53 function reduced its sensitivity to these agents [48].

These *in vitro* studies were complemented by experiments in murine models by Ruzankina et al. [49], who generated ATR-mosaic knockout models from *TP53^{-/-}* mice. Mice with concomitant loss of both ATR and *TP53* displayed markedly reduced survival compared to those harboring isolated loss of ATR and exhibited high levels of DNA damage as evidenced by the accumulation of γ H2AX-positive cells [49]. In a further study using xenotransplantation models of both p53-wild-type and p53-deficient triple-negative breast cancer, Chk1 inhibition was shown to potentiate chemotherapy-induced apoptosis in p53-mutant xenografts but not in p53-wild-type counterparts. Combining Chk1 inhibition with the topoisomerase inhibitor irinotecan resulted in suppression of tumor growth and prolonged survival in xenografts with p53 deficiency but not in wild-type xenografts [50].

In addition to *TP53* loss, cyclin E overexpression has also been reported to exacerbate the effects of ATR pathway inhibition [51]. Cyclin E promotes G1/S cell-cycle progression and hence contributes to a heightened level of cellular replication stress.

10.4.3 Synthetic Lethality With Double-Stranded DNA Repair Defects

Ataxia telangiectasia mutated (ATM) is a kinase that acts upstream of p53 and controls a DDR pathway critical to resolving double-stranded DNA breaks (DSBs). ATM has considerable functional redundancy with the ATR pathway [5,52]. Therefore, a second scenario in which tumor cells can become addicted to the ATR pathway is through ATM loss. Like the *TP53* gene, deletions or mutations of the *ATM* gene are also frequent in various malignancies [53]. Moreover, loss of ATM promotes tumor progression and chemoresistance similar to p53 loss [54–56].

When ATR function is inhibited, stalled replication forks with persistent ssDNA collapse, resulting in the formation of partially replicated sister chromatid fragments with DNA double-stranded ends [5,13,28]. This is a form of DSBs that requires homologous recombination repair (HRR) through the ATM pathway. When HRR is defective, these chromatid fragments accumulate, and while they can be ligated through nonhomologous end joining (NHEJ), the latter is a low-fidelity repair process that potentially gives rise to aberrant sequence deletions or chromosomal translocations. In addition, ATM also plays an important role in cell-cycle regulation through both p53-dependent and -independent mechanisms [57]. Thus, when ATR is inhibited in ATM-deficient cells, accumulation of DNA damage ensues and defective cell-cycle checkpoints due to combined loss of ATR and ATM permit unrestricted entry into mitosis, leading to mitotic catastrophe.

Consistent with this, Reaper et al. [58] showed that ATR inhibition was invariably cytotoxic to ATM or p53-defective tumor cells. Moreover, combination of ATR inhibition with RS-inducing genotoxic agents such as cisplatin and carboplatin was more profoundly synergistic in tumor cells with constitutive ATM or p53 defects, and in cells subjected to ATM pharmacological inhibition or p53 siRNA knockdown [58]. In our recently published study, we extended these observations to primary tumor samples and patient-derived xenotransplantation (PDX) models. ATR inhibition was selectively cytotoxic to primary chronic lymphocytic leukemia (CLL) cells harboring deletions or mutations of *ATM* or *TP53*, in comparison with a panel of CLL samples without these aberrations and with healthy donor peripheral blood lymphocytes. Remarkably, ATR inhibitor treatment in murine xenografts derived from CLL patients with biallelic *ATM* loss led to a reduction not only in tumor load, but also in the proportion of CLL cells with *ATM* defects, indicating specificity of ATR inhibition for these defects [31].

Several other studies have demonstrated synthetic lethality between ATR inhibition and ATM or other HRR defects. ATR inhibition was lethal in cell lines derived from patients suffering from gastric cancer with ATM deficiency, in ovarian cancer with Breast cancer type 1 susceptibility protein 1 (BRCA1) deficiency, and in Rad51-depleted cells [59–62]. Taken together, these studies demonstrate that defects in HRR-dependent DSB repair are markers of sensitivity to ATR pathway inhibition.

Fig. 10.1 summarizes the synthetically lethal interaction between ATR pathway inhibition and ATM or p53 loss.

10.4.4 Other Markers of Sensitivity

A number of additional DNA repair defects have been shown to confer enhanced sensitivity to ATR pathway inhibition. These defects include deficiency in the DNA repair proteins XRCC1, ERCC1, POLD1, or PRIM1 [63–65]. XRCC1 is a protein involved in base excision




FIG. 10.1 A model for synthetic lethality in CLL cells with ATM or p53 deficiency by inhibition of ATR. ATM and ATR are master regulators of DDR, with ATM being activated in response to DNA double-strand breaks, and ATR in response to replication stress. (A) Activation of the ATR pathway leads to cell-cycle arrest mediated primarily

repair and the single-strand break repair pathway, whereas ERCC1 mediates the repair of several types of DNA lesions including bulky adducts, DSBs, and inter-strand cross-links and facilitates the separation of sister chromatids at fragile sites. POLD1 and PRIM1, on the other hand, are involved in DNA replication synthesis. Recently, an RNAi screen identified AT-rich interaction domain 1A (ARID1A) deficiency as an additional marker of sensitivity to ATR inhibition. Mutations in ARID1A are common in cancers and lead to disrupted topoisomerase localization and cell-cycle progression, thus imposing on cancer cells a dependence on the ATR signaling pathway [66].

Finally, a study by Flynn et al. [67] has alluded to the hypersensitivity to ATR inhibition in cancer cells reliant on a mechanism of telomere maintenance known as alternative lengthening of telomeres (ALT), whereby telomeres are elongated through recombination. The absence of ATR leads to abrogation of ALT, compromising telomere stability in ALT-dependent cancer cells, thus resulting in DNA damage, telomere loss, and selective lethality of these cells.

10.5 PRECLINICAL EVALUATION OF SMALL-MOLECULE INHIBITORS TARGETING THE ATR PATHWAY

10.5.1 Development and Evaluation of ATR Pathway Inhibitors

The design and development of selective ATR kinase inhibitors has been coupled with difficulty because of the atypical nature of PIKKs relative to other more conventional kinases. Early inhibitors of ATR were not specific, but more recently two drug companies have manufactured potent and highly specific ATR inhibitors. They are VE-821 and its analog VX-970 produced by Vertex Pharmaceuticals, as well as AZ20 and its analog AZD6738 developed by AstraZeneca [68,69]. Both compounds are ATP-competitive ATR kinase inhibitors, with AZD6738 additionally being available in an oral formulation.

In contrast, Chk1 inhibitors were accessible much earlier, and to date a number of Chk1 inhibitors have been developed. The earlier Chk1 inhibitors, such as UCN-01, however, lacked target selectivity and had poor pharmacokinetic properties, thus limiting their utility [70]. Next-generation Chk1 inhibitors, including AZD7762, MK-8776 (Sch900776), LY2603618, LY2606368, PF-00477736, V158411, and SAR020106, had considerably less off-target effects with an improved pharmacokinetic profile [71].

Numerous preclinical studies on ATR and Chk1 kinase inhibitors have been conducted on an array of solid and hematological malignancies, using a variety of experimental models. Although cell line models of human malignancies are easy to manipulate and provide an excellent basis for the initial evaluation of novel therapeutic agents, effects observed on cell lines

through the G2/M checkpoint, and repair of stalled replication forks. (B–C) Inhibition of ATR leads to collapse of stalled replication forks into partially replicated DNA fragments with double-stranded ends that are repaired through the ATM/p53 pathway. This involves cell-cycle arrest mediated primarily through the G1/S checkpoint and HRR. (D) In cells with defective ATM or p53, inhibition of ATR results in intolerable accumulation of unrepaired DNA damage. This arises from impaired HRR due to defective ATM and/or impaired cell-cycle regulation resulting from combined loss of functional ATR and ATM/p53. Abbreviations: CLL, chronic lymphocytic leukemia; ATM, ataxia telangiectasia mutated; ATR, ataxia telangiectasia and Rad3; HRR, homologous recombination repair. Adapted from Kwok et al., *Blood*, 2016, 127:582–95) [31].

are not always reflective of their effects *in vivo*. Therefore, results from studies on primary tumor material, as well as animal models, especially PDX, are invaluable additions to *in vitro* experimentation in informing the potential drug utility and effects, beneficial or otherwise, prior to their clinical deployment. In the following sections, we examine currently available *in vitro* and *in vivo* preclinical data on ATR and Chk1 inhibitors.

10.5.2 Single-Agent Efficacy of ATR Inhibitors in Cancer

In vitro, exposure of *TP53*-mutant multiple myeloma cell lines to 1 μM of VE-821 for 3 days resulted in a reduction in cell viability by >50% [72]. VE-821 monotherapy was shown to be cytotoxic also to glioma and lymphoma cell lines, to rectal carcinoma cell lines under hypoxic conditions, and to breast cancer and non-small-cell lung cancer lines with ERCC1 defect [60,63,73,74]. In colon cancer, HCT166 cells, particularly those deficient in ARID1A, responded to VE-821, VX-970, and AZ20, as reflected by marked reduction (by >90%) in viability and clonogenic survival upon treatment with ATR inhibitor doses of $\leq 1 \mu\text{M}$ [66]. In acute myeloid leukemia (AML), AZ20 doses of 350 nM to 1.4 μM were sufficient to reduce viability of AML cell lines by 50% (EC_{50}), whereas in primary AML samples this was achieved with AZ20 doses of 800 nM to 27 μM [75]. With AZD6738, single-agent efficacy was seen in cell lines of HER2-positive breast cancer and ATM-deficient gastric cancer [61,76]. Furthermore, in CLL, we demonstrated that AZD6738 treatment for 4 days led to lethality in *TP53* or *ATM* defective CLL cell lines and proliferating primary CLL samples, with an average EC_{50} of 1.4 μM in cell lines and 8.5 μM in primary CLL cells [31].

In vivo, VX-970 suppressed tumor growth and prevented tumor establishment in mice xenografted with ARID1A-deficient HCT116 or ovarian cancer TOV-21G cells, whereas AZ20 showed single-agent efficacy in reducing tumor infiltration and improving survival of murine models transplanted with *MLL*-rearranged AML cells [66,77]. Moreover, AZD6738 monotherapy, administered orally for 2 weeks, led to marked reduction in tumor load in PDX of biallelic *ATM*- or *TP53*-inactivated CLL, accompanied by a reduction in the proportion of CLL cells with these defects [31]. AZD6738 also countered tumor growth and induced apoptosis in a murine xenograft model of ATM-deficient gastric cancer [61].

A summary of the major preclinical studies involving ATR inhibitors in the various cancer types is presented in Table 10.1.

10.5.3 Single-Agent Efficacy of Chk1 Inhibitors in Cancer

In vitro, single-agent Chk1 inhibition with $\leq 1 \mu\text{M}$ AZD7762 led to >90% reduction in the viability of radioresistant breast cancer cell lines as well as viability reductions of varying magnitudes in metastatic melanoma cell lines [78,79]. With MK-8776 (Sch900776), single-agent cytotoxic effects were seen in neuroblastoma lines (median EC_{50} 900 nM), in cell lines of myeloid leukemia and in BRCA-mutant ovarian cancer [80,81]. Cell lines that were sensitive to MK-8776 monotherapy were found to display aberrant Cdk2 activation in S phase upon Chk1 inhibition, leading to DSBs, possibly because these cells depend highly on constitutive suppression of Cdc25 phosphatases by Chk1 [82]. Moreover, LY2603618 exerted single-agent cytotoxicity on AML cell lines (EC_{50} 0.1–1.6 μM) and patient samples (EC_{50} <9 μM) as well as in osteosarcoma cell lines [83,84]. Furthermore, single-agent PF-00477736 activity was evident

TABLE 10.1 Preclinical Studies involving ATR Inhibitors

Malignancy	Study	Inhibitor	Experimental Model	Biomarker of Sensitivity	Treatments Potentiated by ATR Inhibitor
AML	Ma et al., 2017 [75]	AZ20	Cell lines, primary cells	—	Cytarabine
	Chauduri et al., 2014 [93]	VE-821	Cell lines	—	Wee1 inhibitor
Breast	Yazinski et al., 2017 [62]	AZ20, VE-821	Cell lines, primary cells	BRCA1 deficiency	PARP1 inhibitor
	Kim et al., 2017 [76]	AZD6738	Cell lines	—	Cisplatin
Cervical	Teng et al., 2015 [115]	ETP-46464	Cell lines	—	Cisplatin, radiotherapy
CLL	Kwok et al., 2016 [31]	AZD6738	Cell lines, primary cells, PDX	ATM, p53 deficiency	Chlorambucil, fludarabine, cyclophosphamide, bendamustine, ibrutinib
Colorectal	Hocke et al., 2016 [65]	VX-970, NU-6027	Cell lines	POLD1 deficiency, PRIM1 deficiency	—
Endometrial	Teng et al., 2015 [115]	ETP-46464	Cell lines	—	Cisplatin, radiotherapy
Esophageal	Leszczynska et al., 2016 [116]	VX-970	Cell lines, cell-line xenografts	—	Cisplatin, carboplatin, radiotherapy
Gastric	Min et al., 2017 [61]	AZD6738	Cell lines, cell-line xenografts	ATM deficiency	—
Lung	Vendetti et al., 2015 [90]	AZD6738	Cell lines, cell-line xenografts	—	Cisplatin, gemcitabine, radiotherapy
	Hall et al., 2014 [89]	VX-970	Cell lines, primary cells, PDX	—	Cisplatin
Lymphoma	Muralidharan et al., 2016 [73]	AZ20, VE-821	Cell lines, cell-line xenografts	—	BET inhibitor
	Menezes et al., 2015 [117]	WO2010/073034	Cell lines, cell-line xenografts	ATM deficiency	—

(Continued)

TABLE 10.1 Preclinical Studies Involving ATR Inhibitors (*cont.*)

Malignancy	Study	Inhibitor	Experimental Model	Biomarker of Sensitivity	Treatments Potentiated by ATR Inhibitor
Myeloma	Cottini et al., 2015 [72]	VE-821	Cell lines, primary cells	Myc expression, p53 deficiency	Piperlongumine
Ovarian	Kim et al., 2016 [81]	AZD6738	Cell lines, PDX	—	PARP1 inhibitor
	Huntoon et al., 2013 [118]	VE-821	Cell lines	—	Cisplatin, gemcitabine, PARP inhibitor
Pancreatic	Prevo et al., 2012 [119]	VE-821	Cell lines, primary cells	—	Gemcitabine, radiotherapy
	Fokas et al., 2012 [120]	VX-970	Cell lines, cell-line xenografts	—	Gemcitabine, radiotherapy
Various	Toledo et al., 2011 [51]	ETP-46464	Cell lines	p53 deficiency, cyclin E overexpression	Hydroxyurea, Chk1 inhibitor
	Sultana et al., 2013 [64]	NU-6027	Cell lines	XRCC1 deficiency	Cisplatin
	Repear et al., 2011 [58]	VE-821	Cell lines	ATM, p53 deficiency	Cisplatin
	Pires et al., 2012 [74]		Cell lines	—	Radiotherapy
	Mohni et al., 2015 [91]		Cell lines	TLS polymerase, 53BP1	Cisplatin
	Krajewska et al., 2015 [59]		Cell lines	HRR deficient, Rad51 deficiency	—
	Middelton et al., 2015 [60]		Cell lines	HR/BER defects	—
	Josse et al., 2014 [92]	VE-821, VX-970	Cell lines, cell-line xenografts	—	Topoisomerase inhibitor
	Williamson et al., 2016 [66]		Cell lines, cell-line xenografts	ARID1A deficiency	—
	Mohni et al., 2014 [63]	VE-821	Cell lines	ERCC1 deficiency	—

Abbreviations: AML, acute myeloid leukemia; CLL, chronic lymphocytic leukemia; PDX, patient-derived xenografts; TLS, translesion synthesis.

in T-cell acute lymphoblastic leukemia (T-ALL) cell lines (EC_{50} 20–109 nM) and primary cells (EC_{50} <200 nM) which overexpress Chk1 [85]. PF-00477736 also exerted single-agent activity in cell lines of mantle cell lymphoma (EC_{50} 0.68 nM), germinal center B-cell like (GCB) diffuse large B cell lymphoma (DLBCL; EC_{50} 10.2 nM) and activated B-cell (ABC) DLBCL (EC_{50} 87.3 nM) [86]. Finally, V158411 inhibited cell proliferation and induced caspase activation in breast and ovarian cancer cell lines [87].

In vivo, AZD6722 treatment for 3 days resulted in delay in tumor growth in MCF-7/C6 radioresistant breast cancer xenografts [78]. MK-8776 (Sch900776), on the other hand, reduced tumor volume in MDA-MB-231 breast cancer xenografts [88]. Furthermore, in a xenotransplantation model of T-ALL, 30 days of treatment with PF-00477736 yielded a significant reduction in tumor growth compared to vehicle-treated controls [85].

A selection of important preclinical studies in different cancers involving Chk1 inhibitors is presented in Table 10.2.

10.5.4 Chemotherapeutic Combinations Involving ATR Pathway Inhibitors

ATR and Chk1 inhibitors can be readily combined with conventional chemotherapeutic agents. Although the effectiveness of ATR/Chk1 inhibitor monotherapy is often dependent on specific tumor phenotypes, such as p53 deficiency or Myc overexpression, ATR and Chk1 inhibitors, when tested in preclinical models, synergistically enhance the effect of chemotherapy across a broad range of cancer phenotypes, DDR defective or otherwise. This can be ascribed to the potentiation of chemotherapy-induced RS by ATR/Chk1 inhibitors. The ability of ATR pathway inhibitors to resensitize chemoresistant tumor cells to conventional chemotherapeutic agents is of particular clinical significance.

Among the numerous chemotherapeutic agents that synergize with ATR/Chk1 inhibitors, cisplatin and gemcitabine are the most frequently studied. Cisplatin causes DNA breaks and crosslinks, whereas gemcitabine induces RS by decreasing the deoxyribonucleotide triphosphate required for DNA replication. ATR inhibitors broadly sensitized a panel of lung cancer cell lines, cell line xenografts, and PDX to cisplatin or gemcitabine, and ovarian cancer cell lines to cisplatin [89,90]. The sensitization to cisplatin is especially profound in tumor cells with deficiency in ATM, p53, or XRCC1, or with loss of 53BP1 or polymerase ζ involved in translesion synthesis [48,58,64,91]. Other DNA damaging agents that synergize with ATR inhibitors include cytarabine, as observed in AML cell lines and primary cells, as well as chlorambucil, bendamustine, fludarabine, and cyclophosphamide, as observed in CLL cell lines and primary cells. ATR inhibitors also sensitized tumor cells to topoisomerase inhibitors through disruption of DNA replication initiation and fork elongation processes [92].

Furthermore, ATR pathway inhibitors display synergy across a range of novel agents. Two studies, for instance, have highlighted the synergistic interaction between poly-(ADP-ribose) polymerase (PARP) inhibitors and ATR or Chk1 inhibitors in ovarian and breast cancer, respectively, using cell lines complemented by primary tumor samples or PDX [62,81]. PARP inhibitor treatment led to increased accumulation of cells in G2 phase, thus intensifying their reliance on the ATR pathway for checkpoint control and the maintenance of genome stability. ATR and/or Chk1 inhibitors have also shown synergy with inhibitors of other DNA repair molecules, such as Wee1, which contributes to the regulation of the G2/M checkpoint, and MK2, which is critical for prolonged checkpoint maintenance [79,86,93–96]. Finally,

TABLE 10.2 Preclinical Studies Involving Chk1 Inhibitors

Malignancy	Study	Inhibitor	Experimental Model	Biomarker of Sensitivity	Treatments Potentiated by Chk1 Inhibitor
ALL	Sarmiento et al., 2015 [85]	PF477736	Cell lines, cell-line xenografts, primary cells	—	—
AML	Zhao et al., 2016 [83]	LY2603618	Cell lines, primary cells	—	Bcl-2 inhibitor
	Chauduri et al., 2014 [93]	MK-8776	Cell lines	—	Wee1 inhibitor
	Yuan et al., 2014 [80]	SCH900766, AZD7762	Cell lines	—	—
Bladder	Wang et al., 2015 [121]	Gö6976	Cell lines	—	Gemcitabine
Breast	Zhang et al., 2016 [78]	AZD7762	Cell lines, cell-line xenografts	—	—
	Zhou et al., 2017 [88]	MK-8776	Cell lines, cell-line xenografts	—	Radiotherapy
	Ma et al., 2012 [50]	UCN-01, AZD7762	PDX	p53 deficiency	Topoisomerase inhibitor
	Tang et al., 2012 [122]		Cell lines, cell-line xenografts	—	PARP inhibitor, radiotherapy
	Bryant et al., 2014 [87]	V158411, PF477736, AZD7762	Cell lines	—	Gemcitabine, cisplatin
Colon	Martino-Echarri et al., 2014 [123]	MK-8776, AZD7762	Cell lines	—	5-Fluorouracil
	Origanti et al., 2013 [124]	UCN-01	Murine models, orthotopic models	p53 deficiency, p21 deficiency	Topoisomerase inhibitor
Head & neck	Gadhikar et al., 2013 [125]	AZD7762	Cell lines	p53 deficiency	Cisplatin
	Barker et al., 2016 [126]	CCT244747	Cell lines, cell-line xenografts, primary cells	—	Paclitaxel
Lung	Bartucci et al., 2012 [127]	SB218078, AZD7762	Primary cells, PDX	—	Gemcitabine, cisplatin, paclitaxel

TABLE 10.2 Preclinical Studies Involving Chk1 Inhibitors (*cont.*)

Malignancy	Study	Inhibitor	Experimental Model	Biomarker of Sensitivity	Treatments Potentiated by Chk1 Inhibitor
Lymphoma	Chila et al., 2015 [86]	PF477736	Cell lines, cell-line xenografts	—	Wee1 inhibitor
	Zemanova et al., 2016 [128]	SCH900776	Cell lines, primary cells, murine models	p53 deficiency	Fludarabine, cytarabine, gemcitabine
	Murga et al., 2011 [42]	UCN-01	Murine models	Myc expression	—
Melanoma	Magnussen et al., 2015 [79]	AZD7762	Cell lines, cell-line xenografts	—	Wee1 inhibitor
Myeloma	Pei et al., 2014 [129]	CEP3891	Cell lines, primary cells	—	MEK1/2 inhibitor
	Dai et al., 2011 [130]	UCN-01	Cell lines, primary cells, PDX	—	SRC inhibitor
Nasopharyngeal	Mak et al., 2015 [95]	AZD7762	Cell lines, cell-line xenografts	—	Wee1 inhibitor
Neuroblastoma	Russell et al., 2013 [94]	MK-8776	Cell lines, cell-line xenografts	—	Wee1 inhibitor
	Cole et al., 2011 [131]	SB218078, TCS2312	Cell lines, primary cells	Myc expression	—
Oral SCC	Sankunny et al., 2014 [132]	PF477736	Cell lines	ATM deficiency	Radiotherapy
Osteosarcoma	Duan et al., 2014 [84]	LY2603618	Cell lines	—	Cisplatin
Ovarian	Kim et al., 2016 [81]	MK-8776	Cell lines, PDX	—	PARP inhibitor
	Bryant et al., 2014 [87]	V158411, PF477736, AZD7762	Cell lines	—	Gemcitabine, cisplatin
Pancreatic	Morgan et al., 2010 [92]	AZD7762	Cell lines, cell-line xenografts, PDX	—	Radiotherapy
	Vance et al., 2011 [133]		Cell lines	p53 deficiency	PARP inhibitor
	Engelke et al., 2013 [134]	MK-8776	Cell lines, cell-line xenografts	HRR deficiency	Gemcitabine, radiotherapy

(Continued)

TABLE 10.2 Preclinical Studies Involving Chk1 Inhibitors (*cont.*)

Malignancy	Study	Inhibitor	Experimental Model	Biomarker of Sensitivity	Treatments Potentiated by Chk1 Inhibitor
Various	McNeely et al., 2010 [135]	AZD7762	Cell lines	BRCA2 deficiency, XRCC3 deficiency, DNA-PK deficiency	Gemcitabine
	Krajewska et al., 2015 [59]		Cell lines	HRR deficient, Rad51 deficiency	—
	Sakurikar et al., 2016 [82]	MK-8776	Cell lines	CDK2 activation in S phase	—
	Dietlein et al., 2015 [96]	PF477736	Cell lines, cell-line xenografts, primary cells	KRAS mutation	MK2 inhibitor

Abbreviations: ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; SCC, squamous cell carcinoma; PDX, patient-derived xenografts.

therapeutic combinations with ATR pathway inhibitors are not confined to agents targeting DDR. ATR inhibitors are capable of enhancing the effect of a BET bromodomain inhibitor in Myc-driven lymphoma and a B-cell receptor signaling inhibitor in CLL [31,73].

10.6 CLINICAL TRIALS OF SMALL-MOLECULE INHIBITORS TARGETING THE ATR PATHWAY

10.6.1 Clinical Studies Involving Chk1 Inhibitors

To explore the use of selective Chk1 inhibitors in oncology practice, several clinical studies have been carried out on AZD7762, LY2603618, LY2606368, and MK-8776 (Sch900776). These were phase I or II trials with the aim to assess safety and tolerability, to establish the maximum tolerable dose and, within certain studies, to determine response rate and durability. Unfortunately, the clinical development of AZD7762 was hampered by drug-related cardiac toxicity, which was dose limiting in some cases [97,98]. In contrast, LY2603618, LY2606368, and MK-8776 (Sch900776) had an acceptable toxicity profile with rare serious adverse events [99–102]. Moreover, early evidence of clinical activity was seen with these agents. In particular, eight (out of 24) refractory AML patients achieved complete remission and 15 (out of 30) individuals with advanced solid malignancy had partial response or stable disease following treatment with MK-8776 (Sch900776), either alone or in combination with chemotherapy [101,102]. Likewise, partial response was seen in two patients with advanced cancer following LY2606368 monotherapy [100].

Despite this, two phase II trials investigating LY2603618 in advanced lung and pancreatic cancer reported nonsuperiority of LY2603618, used in combination with premetrexed or gemcitabine, respectively, when compared to premetrexed or gemcitabine alone [103,104]. However, since the patients enrolled on these trials were unselected, a substantial proportion would unlikely have harbored high RS phenotypes or lesions that would yield synthetic lethality with ATR/Chk1 inhibition. Indeed, genomic analysis of an exceptional responder, who was cured of her invasive ureteric cancer with AZD7762 in combination with irinotecan, revealed mutation within *RAD50* which attenuated ATM signaling and enforced dependence on the ATR pathway [105]. This underscores the importance of the use of ATR pathway inhibitors for individuals who are likely to benefit.

Unfortunately, the clinical development of AZD7762, LY2603618, and MK-8776 has been suspended. Currently, the Chk1 inhibitors under active clinical investigation include LY2606368 and CCT245737, as detailed in Table 10.3.

10.6.2 Clinical Studies Involving ATR Inhibitors

With regard to ATR inhibitors, both AZD6738 and VX-970 have now entered phase I/II clinical testing. However, the majority of these studies have been initiated recently, and no results have yet been reported. As of March 2017, 11 clinical trials on ATR inhibitors are ongoing for a range of malignancies. These studies examine the use of the ATR inhibitor either alone or in combination with a range of conventional and novel therapeutic agents, including cisplatin, carboplatin, gemcitabine, etoposide, irinotecan, the PARP inhibitors olaparib and veliparib, and the anti-PD-L1 immune checkpoint inhibitor durvalumab.

A summary of the completed and ongoing clinical trials involving Chk1 and ATR inhibitors is presented in Table 10.3.

10.7 TARGETING THE ATM PATHWAY: IS THIS OF ANY VALUE IN OVERCOMING CHEMORESISTANCE IN CANCER?

There is considerable cross-talk between the ATR and ATM signaling pathways. Given the central importance of ATM in DDR, it is reasonable to also examine whether there is potential utility in inhibiting ATM to overcome chemoresistance, particularly since a number of selective ATM inhibitors are now available, including KU-55933, KU-60019, KU-59403, and AZD0156. In comparison with the plethora of literature on ATR pathway targeting, the number of studies on ATM targeting is relatively limited. It is generally recognized that ATM-deficient cells are hypersensitive to ionizing radiation (IR) on account of their inability to repair DSBs. Therefore, historical studies have focused on the use of ATM inhibitors to sensitize tumor cells to radiotherapy. Indeed, ATM inhibitors are potent radiosensitizers, both *in vitro* and *in vivo* [106–108].

More recently, Batey et al. [109] showed that ATM inhibitors can also be used to potentiate the cytotoxic effect of topoisomerase inhibitors such as camptothecin, etoposide, and doxorubicin, but lacked single-agent activity. However, the prolonged use of ATM inhibition remains questionable. In contrast to ATR inhibitors with target predilection for p53- and HRR-deficient cells, the lack of such predilection of ATM inhibitors raises the possibility of toxicity

TABLE 10.3 Clinical Studies Involving Chk1 and ATR Inhibitors

Inhibitor	Phase	Study	Treatment	Malignancy	Pt no	Endpoint	Conclusions
AZD7762	I	Seto et al., 2013 [98]	Monotherapy; combination with gemcitabine	Advanced solid tumor	20	Safety, pharmacokinetics, preliminary efficacy	Cardiac and hematologic toxicity was dose limiting at 30 mg. Maximum tolerable dose of AZD7762 in combination with gemcitabine was 21 mg
		Sausville et al., 2014 [97]	Monotherapy; combination with gemcitabine	Advanced solid tumor	42	Safety, tolerability	Cardiac toxicity was dose limiting
LY2603618	I	Doi et al., 2015 [99]	Monotherapy; combination with gemcitabine	Advanced solid tumor	17	Safety, tolerability	Acceptable safety and tolerability in combination with gemcitabine. Partial response in four patients. Most common high-grade toxicities were hematological
		Calvo et al., 2016 [136]	Monotherapy; combination with gemcitabine	Advanced/metastatic solid tumor	50	Safety, tolerability	Adverse events include fatigue, thrombocytopenia, neutropenia, nausea, and anemia. Maximum tolerated dose was 200 mg/m ² when combined with gemcitabine. Proceed to phase II trial with a fixed dose of 230 mg
	II	Scagliotti et al., 2016 [103]	Combination with pemetrexed	Metastatic non-small cell lung cancer	53	Response rate, pharmacokinetics, safety	Partial response in five patients. No association between p53 status and response. Efficacy and safety similar to pemetrexed monotherapy
		Laguerre et al., 2017 [104]	Combination with gemcitabine	Metastatic pancreatic cancer	99	Response rate, PFS, pharmacokinetics, safety	Maximum pharmacodynamic response was achieved after 230 mg of LY2603618. Combination of gemcitabine with LY2603618 was not superior to gemcitabine alone

LY2606368	I	Hong et al., 2016 [100]	Monotherapy	Advanced cancer	45	Safety, pharmacokinetics, pharmacodynamics	Dose-limiting toxicities in seven patients were all hematological. Partial response was seen in two patients. Phase II study will utilize LY2606368 dose of 105 mg/m ²
MK-8776 (Sch900776)	I	Karp et al., 2012 [101]	Combination with cytarabine	Refractory acute leukemia	24	Safety, tolerability, preliminary efficacy	Complete response seen in eight patients (33%). Dose-limiting toxicities were cardiac. Recommended dose of Sch900776 for phase II study is 100 mg
		Daud et al., 2015 [102]	Monotherapy; combination with gemcitabine	Advanced solid tumor	43	Safety, pharmacokinetics, pharmacodynamics	Well tolerated as single agent and in combination with gemcitabine. Of 30 evaluable patients, 2 showed partial response and 13 had stable disease. Recommended phase II dose was 200 mg MK-8776 + 1000 mg/m ² gemcitabine

(Continued)

TABLE 10.3 Clinical Studies Involving Chk1 and ATR Inhibitors (cont.)

Inhibitor	Phase	Start Date	Malignancy	Treatment	Pt no	ClinicalTrials.gov Identifier
ATR	I	2014	Advanced solid tumor	Monotherapy; combination with palliative radiotherapy	100	NCT02223923
			Advanced solid tumor with ATM deficiency	Monotherapy; combination with carboplatin, olaparib, or durvalumab	114	NCT02264678
VX-970	I	2015	Advanced solid tumor	Combination with paclitaxel	21	NCT02630199
		2012	Advanced solid tumor	Combination with gemcitabine, cisplatin, etoposide, and/or carboplatin	205	NCT02157792
		2016	Advanced head and neck squamous cell carcinoma	Combination with cisplatin/X-ray therapy	45	NCT02567422
			Non-small cell lung cancer with brain metastases	Combination with whole brain radiotherapy	46	NCT02589522
			Advanced solid tumor	Combination with irinotecan	51	NCT02595931
			Advanced solid tumor	Combination with veliparib and cisplatin	60	NCT02723864
	I/II	2015	Small cell lung cancer	Combination with topotecan	55	NCT02487095
		2016	Metastatic urothelial cancer	Combination with cisplatin/gemcitabine (vs cisplatin/gemcitabine alone)	90	NCT02567409
Chk1	I	2016	Platinum-resistant recurrent ovarian or fallopian tube cancer	Combination with gemcitabine (vs gemcitabine alone)	70	NCT02595892
			Advanced solid tumor	Monotherapy	40	NCT02797964
LY 2606368	I	2017	Advanced solid tumor	Monotherapy	70	NCT02797977
	I	2017	Advanced pediatric solid tumors	Monotherapy	65	NCT02808650
	II	2014	BRC-A mutant or high-grade breast and ovarian cancer, metastatic prostate cancer	Monotherapy	144	NCT02203513
			Solid tumors with replication stress or HRR deficiency	Monotherapy	38	NCT02873975

Published studies are listed in the upper panel, and ongoing studies in the lower panel. Abbreviations: Pt no, patient number; PFS, progression-free survival.

to healthy cells [109]. ATM inhibitors may be more effective in certain tumor types, such as glioblastoma, where tumor-associated overexpression of platelet-derived growth factor alpha (PDGFRA) can be normalized with ATM inhibition [110]. While there is no published clinical data on ATM inhibition, a phase I study of AZD0156, alone and in combination with chemotherapy or olaparib, is currently underway (NCT02588105). Therefore, the clinical utility of ATM inhibition remains to be seen.

10.8 CONSIDERATIONS, CAVEATS, AND UNRESOLVED QUESTIONS IN ATR TARGETING

10.8.1 ATR vs Chk1 targeting

While ATR and Chk1 act in the same pathway, ATR is upstream of Chk1 and therefore controls additional downstream processes. Targeting ATR may therefore produce more wide-ranging effects than targeting Chk1, and this is reflected in the ability of ATR inhibitors to potentiate a broader range of chemotherapeutic agents. It remains to be determined whether ATR inhibitors are clinically more effective than Chk1 inhibitors. This might vary from one malignancy to another, or indeed from one tumor phenotype to another. For example, tumors with ATM deficiency might benefit more from ATR inhibition than from Chk1 inhibition, given the functional cross-talk between the ATM and ATR.

10.8.2 Potential Mechanisms of Resistance to ATR Pathway Inhibition

A fundamental assumption underpinning models of synthetic lethality is that there are only two major pathways regulating a process. Therefore, if one pathway is defective, cellular demise is assured when the other pathway is blocked. However, this notion is inevitably an oversimplification of the myriad of collateral pathways regulating a cellular process, many of which are hitherto underappreciated or unknown. When one collateral pathway is therapeutically inhibited, tumor cells may upregulate an alternative collateral pathway to mitigate the effects of the initial block, thereby resulting in therapeutic resistance.

Alongside ATM, DNA-dependent protein kinase (DNA-PK) plays a role in DSB repair through control of NHEJ. Recently, there have been reports pointing to a possible functional redundancy between ATR and DNA-PK in regulating downstream Chk1 activity [111]. Buisson et al. [112], in particular, demonstrated the existence of a DNA-PK-Chk1 backup pathway that can mediate resistance to ATR inhibitors. In this model, ATR inhibition is cytotoxic to a proportion of tumor cells with the highest levels of replication stress, but those with moderate levels are protected through the backup pathway [112]. Simultaneous targeting of ATR and Chk1 or DNA-PK could potentially overcome this. Indeed, a potentiating interaction between ATR and Chk1 inhibitors has been reported, suggesting that the combined use of these inhibitors could be more efficacious than either agent alone [113]. However, the potential toxicities and adverse effects of any combined use of ATR and Chk1 inhibitors will need to be properly assessed.

Finally, both ATR and Chk1 inhibitors are kinase inhibitors. As observed from the small-molecule kinase inhibitors currently in clinical use, point mutations in the target kinase can

develop. This constitutes another potential source of therapeutic resistance that physicians and investigators will need to anticipate and overcome.

10.8.3 Precision Medicine and the Selection of Patients for ATR/Chk1 Targeting

The long-term effects and toxicity of ATR and Chk1 inhibition are unknown and this is often a source of contention. Targeting a process of such importance to genome integrity can be potentially dangerous if it also affects healthy cells. Moreover, sublethal targeting of tumor cells with ATR pathway inhibitors can inadvertently promote tumorigenesis and clonal evolution, as these cells accumulate replication stress and DNA damage but escape death [114]. It is therefore of paramount importance that patients are selected for ATR pathway inhibitor treatment according to whether their tumor cells are differentially sensitive to these inhibitors. By selecting patients whose tumor is hypersensitive to these inhibitors, tumor killing can be maximized while minimizing treatment duration, thereby reducing both toxicity to healthy tissues and the risk of tumor evolution and escape. While a number of biomarkers of sensitivity to ATR pathway inhibition have been discussed earlier, it is likely that many others have yet to be discovered. The onus is on investigators to continue to identify predictive biomarkers of sensitivity to ATR and Chk1 inhibitors, specific for each tumor type, in order to enable accurate therapeutic stratification.

10.8.4 The Use of ATR/Chk1 Inhibitors with Other Therapeutic Agents

Finally, should ATR/Chk1 inhibitors be administered as a single agent or in combination with chemotherapy? If so, what is the ideal therapeutic regimen to be used? An argument in favor of combined use with chemotherapy is that the addition of the latter would increase the potency of ATR pathway inhibitors, but this needs to be balanced by a possible increase in toxicity. In addition, the use of therapeutic combinations may potentially reduce the likelihood of developing resistance to single agents. The ideal chemotherapeutic agent to be used in combination with different tumor types will need to be ascertained through mechanistic and clinical studies. In addition, ATR/Chk1 inhibitors can potentially be combined with other novel small-molecule inhibitors or monoclonal antibodies. This may be of particular relevance to a number of malignancies where a substantial proportion of tumor cells is quiescent. These cells are unlikely to be susceptible to ATR pathway inhibition and may give rise to a pool of residual cells from which therapeutic resistance can arise. Therefore, combination of ATR pathway inhibitors with agents targeting pathways unrelated to replication could allow the simultaneous eradication of both proliferating and quiescent tumor populations. The scheduling and sequencing of ATR pathway inhibitors within therapeutic regimens will require continuous optimization.

10.8.5 Concluding Remarks

The availability of ATR pathway inhibitors offer a beacon of hope that chemoresistance, which is a barrier to effective cancer treatment, could finally be overcome. Preclinical results

of ATR and Chk1 inhibitors have shown remarkable promise. However, our understanding of the mechanistic basis of ATR pathway inhibition is by no means complete, and the clinical development of ATR pathway inhibitors is still in its infancy. Identification of predictive biomarkers of response underpinned by sound mechanistic understanding remains a research priority, and the rational selection of patients into clinical studies based on these biomarkers could translate into better therapeutic outcomes. In conclusion, the development of ATR pathway inhibitors epitomizes the emerging role of precision medicine in cancer treatment, through which vulnerabilities in cancer are systematically identified and targeted.

Acknowledgments

We would like to acknowledge Bloodwise UK for continuous support of our studies.

Conflict of Interest: No potential conflicts of interest were disclosed.

References

- [1] Jackson SP, Bartek J. The DNA-damage response in human biology and disease. *Nature* 2009;461(7267):1071–8.
- [2] Castedo M, Perfettini JL, Roumier T, Andreau K, Medema R, Kroemer G. Cell death by mitotic catastrophe: a molecular definition. *Oncogene* 2004;23(16):2825–37.
- [3] Vitale I, Galluzzi L, Castedo M, Kroemer G. Mitotic catastrophe: a mechanism for avoiding genomic instability. *Nat Rev Mol Cell Biol* 2011;12(6):385–92.
- [4] O'Connor MJ. Targeting the DNA damage response in cancer. *Mol Cell* 2015;60(4):547–60.
- [5] Cimprich KA, Cortez D. A.T.R.: an essential regulator of genome integrity. *Nat Rev Mol Cell Biol* 2008;9(8):616–27.
- [6] Ge XQ, Blow JJ. Chk1 inhibits replication factory activation but allows dormant origin firing in existing factories. *J Cell Biol* 2010;191(7):1285–97.
- [7] Chen YH, Jones MJ, Yin Y, et al. ATR-mediated phosphorylation of FANCI regulates dormant origin firing in response to replication stress. *Mol Cell* 2015;58(2):323–38.
- [8] Marechal A, Zou L. RPA-coated single-stranded DNA as a platform for post-translational modifications in the DNA damage response. *Cell Res* 2015;25(1):9–23.
- [9] Cotta-Ramusino C, McDonald ER III, Hurov K, Sowa ME, Harper JW, Elledge SJ. A DNA damage response screen identifies RHINO, a 9-1-1 and TopBP1 interacting protein required for ATR signaling. *Science* 2011;332(6035):1313–7.
- [10] Haahr P, Hoffmann S, Tollenaere MA, et al. Activation of the ATR kinase by the RPA-binding protein ETAA1. *Nat Cell Biol* 2016;18(11):1196–207.
- [11] Bass TE, Luzwick JW, Kavanaugh G, et al. ETAA1 acts at stalled replication forks to maintain genome integrity. *Nat Cell Biol* 2016;18(11):1185–95.
- [12] Lee YC, Zhou Q, Chen J, Yuan J. RPA-binding protein ETAA1 is an ATR activator involved in DNA replication stress response. *Curr Biol* 2016;26(24):3257–68.
- [13] Dungrawala H, Rose KL, Bhat KP, et al. The replication checkpoint prevents two types of fork collapse without regulating replisome stability. *Mol Cell* 2015;59(6):998–1010.
- [14] Neelsen KJ, Lopes M. Replication fork reversal in eukaryotes: from dead end to dynamic response. *Nat Rev Mol Cell Biol* 2015;16(4):207–20.
- [15] Berti M, Vindigni A. Replication stress: getting back on track. *Nat Struct Mol Biol* 2016;23(2):103–9.
- [16] Ammazalorso F, Pirzio LM, Bignami M, Franchitto A, Pichierri P. ATR and ATM differently regulate WRN to prevent DSBs at stalled replication forks and promote replication fork recovery. *EMBO J* 2010;29(18):3156–69.
- [17] Thangavel S, Berti M, Levikova M, et al. DNA2 drives processing and restart of reversed replication forks in human cells. *J Cell Biol* 2015;208(5):545–62.
- [18] Couch FB, Bansbach CE, Driscoll R, et al. ATR phosphorylates SMARCAL1 to prevent replication fork collapse. *Genes Dev* 2013;27(14):1610–23.
- [19] Betous R, Mason AC, Rambo RP, et al. SMARCAL1 catalyzes fork regression and Holliday junction migration to maintain genome stability during DNA replication. *Genes Dev* 2012;26(2):151–62.

- [20] Singh TR, Ali AM, Paramasivam M, et al. ATR-dependent phosphorylation of FANCM at serine 1045 is essential for FANCM functions. *Cancer Res* 2013;73(14):4300–10.
- [21] Schwab RA, Blackford AN, Niedzwiedz W. ATR activation and replication fork restart are defective in FANCM-deficient cells. *EMBO J* 2010;29(4):806–18.
- [22] Davies SL, North PS, Hickson ID. Role for BLM in replication-fork restart and suppression of origin firing after replicative stress. *Nat Struct Mol Biol* 2007;14(7):677–9.
- [23] Iannascoli C, Palermo V, Murfuni I, Franchitto A, Pichierri P. The WRN exonuclease domain protects nascent strands from pathological MRE11/EXO1-dependent degradation. *Nucleic Acids Res* 2015;43(20):9788–803.
- [24] Petermann E, Woodcock M, Helleday T. Chk1 promotes replication fork progression by controlling replication initiation. *Proc Natl Acad Sci USA* 2010;107(37):16090–5.
- [25] Fragkos M, Ganier O, Coulombe P, Mechali M. DNA replication origin activation in space and time. *Nat Rev Mol Cell Biol* 2015;16(6):360–74.
- [26] Peng CY, Graves PR, Thoma RS, Wu Z, Shaw AS, Piwnicka-Worms H. Mitotic and G2 checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on serine-216. *Science* 1997;277(5331):1501–5.
- [27] Sorensen CS, Syljuasen RG, Falck J, et al. Chk1 regulates the S phase checkpoint by coupling the physiological turnover and ionizing radiation-induced accelerated proteolysis of Cdc25A. *Cancer Cell* 2003;3(3):247–58.
- [28] Zeman MK, Cimprich KA. Causes and consequences of replication stress. *Nat Cell Biol* 2014;16(1):2–9.
- [29] Mankouri HW, Huttner D, Hickson ID. How unfinished business from S-phase affects mitosis and beyond. *EMBO J* 2013;32(20):2661–71.
- [30] Toledo LI, Altmeyer M, Rask MB, et al. ATR prohibits replication catastrophe by preventing global exhaustion of RPA. *Cell* 2013;155(5):1088–103.
- [31] Kwok M, Davies N, Agathangelou A, et al. ATR inhibition induces synthetic lethality and overcomes chemoresistance in TP53- or ATM-defective chronic lymphocytic leukemia cells. *Blood* 2016;127(5):582–95.
- [32] Macheret M, Halazonetis TD. DNA replication stress as a hallmark of cancer. *Annu Rev Pathol* 2015;10:425–48.
- [33] Hills SA, Diffley JF. DNA replication and oncogene-induced replicative stress. *Curr Biol* 2014;24(10):R435–44.
- [34] Brown EJ, Baltimore D. Essential and dispensable roles of ATR in cell cycle arrest and genome maintenance. *Genes Dev* 2003;17(5):615–28.
- [35] Zigelboim I, Schmidt AP, Gao F, et al. ATR mutation in endometrioid endometrial cancer is associated with poor clinical outcomes. *J Clin Oncol* 2009;27(19):3091–6.
- [36] Chen CF, Ruiz-Vega R, Vasudeva P, et al. ATR mutations promote the growth of melanoma tumors by modulating the immune microenvironment. *Cell Rep* 2017;18(10):2331–42.
- [37] de Klein A, Muijtjens M, van Os R, et al. Targeted disruption of the cell-cycle checkpoint gene ATR leads to early embryonic lethality in mice. *Curr Biol* 2000;10(8):479–82.
- [38] Ruzankina Y, Pinzon-Guzman C, Asare A, et al. Deletion of the developmentally essential gene ATR in adult mice leads to age-related phenotypes and stem cell loss. *Cell Stem Cell* 2007;1(1):113–26.
- [39] O'Driscoll M, Ruiz-Perez VL, Woods CG, Jeggo PA, Goodship JA. A splicing mutation affecting expression of ataxia-telangiectasia and Rad3-related protein (ATR) results in Seckel syndrome. *Nat Genet* 2003;33(4):497–501.
- [40] Murga M, Bunting S, Montana MF, et al. A mouse model of ATR-Seckel shows embryonic replicative stress and accelerated aging. *Nat Genet* 2009;41(8):891–8.
- [41] Schoppy DW, Ragland RL, Gilad O, et al. Oncogenic stress sensitizes murine cancers to hypomorphic suppression of ATR. *J Clin Invest* 2012;122(1):241–52.
- [42] Murga M, Campaner S, Lopez-Contreras AJ, et al. Exploiting oncogene-induced replicative stress for the selective killing of Myc-driven tumors. *Nat Struct Mol Biol* 2011;18(12):1331–5.
- [43] McLornan DP, List A, Mufti GJ. Applying synthetic lethality for the selective targeting of cancer. *N Engl J Med* 2014;371(18):1725–35.
- [44] Shaheen M, Allen C, Nickoloff JA, Hromas R. Synthetic lethality: exploiting the addiction of cancer to DNA repair. *Blood* 2011;117(23):6074–82.
- [45] Menendez D, Inga A, Resnick MA. The expanding universe of p53 targets. *Nat Rev Cancer* 2009;9(10):724–37.
- [46] Muller PA, Vousden KH. p53 mutations in cancer. *Nat Cell Biol* 2013;15(1):2–8.
- [47] Nghiem P, Park PK, Kim Y, Vaziri C, Schreiber SL. ATR inhibition selectively sensitizes G1 checkpoint-deficient cells to lethal premature chromatin condensation. *Proc Natl Acad Sci USA* 2001;98(16):9092–7.
- [48] Sangster-Guity N, Conrad BH, Papadopoulos N, Bunz F. ATR mediates cisplatin resistance in a p53 genotype-specific manner. *Oncogene* 2011;30(22):2526–33.

- [49] Ruzankina Y, Schoppy DW, Asare A, Clark CE, Vonderheide RH, Brown EJ. Tissue regenerative delays and synthetic lethality in adult mice after combined deletion of Atr and Trp53. *Nat Genet* 2009;41(10):1144–9.
- [50] Ma CX, Cai S, Li S, et al. Targeting Chk1 in p53-deficient triple-negative breast cancer is therapeutically beneficial in human-in-mouse tumor models. *J Clin Invest* 2012;122(4):1541–52.
- [51] Toledo LI, Murga M, Zur R, et al. A cell-based screen identifies ATR inhibitors with synthetic lethal properties for cancer-associated mutations. *Nat Struct Mol Biol* 2011;18(6):721–7.
- [52] Marechal A, Zou L. DNA damage sensing by the ATM and ATR kinases. *Cold Spring Harb Perspect Biol* 2013;5(9). a012716.
- [53] Choi M, Kipps T, Kurzrock R. ATM mutations in cancer: therapeutic implications. *Mol Cancer Ther* 2016;15(8):1781–91.
- [54] Austen B, Powell JE, Alvi A, et al. Mutations in the ATM gene lead to impaired overall and treatment-free survival that is independent of IGVH mutation status in patients with B-CLL. *Blood* 2005;106(9):3175–82.
- [55] Austen B, Skowronska A, Baker C, et al. Mutation status of the residual ATM allele is an important determinant of the cellular response to chemotherapy and survival in patients with chronic lymphocytic leukemia containing an 11q deletion. *J Clin Oncol* 2007;25(34):5448–57.
- [56] Skowronska A, Parker A, Ahmed G, et al. Biallelic ATM inactivation significantly reduces survival in patients treated on the United Kingdom Leukemia Research Fund Chronic Lymphocytic Leukemia 4 trial. *J Clin Oncol* 2012;30(36):4524–32.
- [57] Shiloh Y, Ziv Y. The ATM protein kinase: regulating the cellular response to genotoxic stress, and more. *Nat Rev Mol Cell Biol* 2013;14(4):197–210.
- [58] Reaper PM, Griffiths MR, Long JM, et al. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. *Nat Chem Biol* 2011;7(7):428–30.
- [59] Krajewska M, Fehrmann RS, Schoonen PM, et al. ATR inhibition preferentially targets homologous recombination-deficient tumor cells. *Oncogene* 2015;34(26):3474–81.
- [60] Middleton FK, Patterson MJ, Elstob CJ, et al. Common cancer-associated imbalances in the DNA damage response confer sensitivity to single agent ATR inhibition. *Oncotarget* 2015;6(32):32396–409.
- [61] Min A, Im SA, Jang H, et al. AZD6738, a novel oral inhibitor of ATR, induces synthetic lethality with ATM-deficiency in gastric cancer cells. *Mol Cancer Ther* 2017;16(4):566–77.
- [62] Yazinski SA, Comaills V, Buisson R, et al. ATR inhibition disrupts rewired homologous recombination and fork protection pathways in PARP inhibitor-resistant BRCA-deficient cancer cells. *Genes Dev* 2017;31(3):318–32.
- [63] Mohni KN, Kavanaugh GM, Cortez D. ATR pathway inhibition is synthetically lethal in cancer cells with ERCC1 deficiency. *Cancer Res* 2014;74(10):2835–45.
- [64] Sultana R, Abdel-Fatah T, Perry C, et al. Ataxia telangiectasia mutated and Rad3 related (ATR) protein kinase inhibition is synthetically lethal in XRCC1 deficient ovarian cancer cells. *PLoS ONE* 2013;8(2):e57098.
- [65] Hocke S, Guo Y, Job A, et al. A synthetic lethal screen identifies ATR-inhibition as a novel therapeutic approach for POLD1-deficient cancers. *Oncotarget* 2016;7(6):7080–95.
- [66] Williamson CT, Miller R, Pemberton HN, et al. ATR inhibitors as a synthetic lethal therapy for tumours deficient in ARID1A. *Nat Commun* 2016;7:13837.
- [67] Flynn RL, Cox KE, Jeitany M, et al. Alternative lengthening of telomeres renders cancer cells hypersensitive to ATR inhibitors. *Science* 2015;347(6219):273–7.
- [68] Charrier JD, Durrant SJ, Golec JM, et al. Discovery of potent and selective inhibitors of ataxia telangiectasia mutated and Rad3 related (ATR) protein kinase as potential anticancer agents. *J Med Chem* 2011;54(7):2320–30.
- [69] Foote KM, Blades K, Cronin A, et al. Discovery of 4-[4-[(3R)-3-methylmorpholin-4-yl]-6-[1-(methylsulfonyl)cyclopropyl]pyrimidin-2-yl]-1H-indole (AZ20): a potent and selective inhibitor of ATR protein kinase with monotherapy *in vivo* antitumor activity. *J Med Chem* 2013;56(5):2125–38.
- [70] McNeely S, Beckmann R, Bence Lin AK. CHEK again: revisiting the development of CHK1 inhibitors for cancer therapy. *Pharmacol Ther* 2014;142(1):1–10.
- [71] Ma CX, Janetka JW, Piwnicka-Worms H. Death by releasing the breaks: CHK1 inhibitors as cancer therapeutics. *Trends Mol Med* 2011;17(2):88–96.
- [72] Cottini F, Hideshima T, Suzuki R, et al. Synthetic lethal approaches exploiting DNA damage in aggressive myeloma. *Cancer Discov* 2015;5(9):972–87.
- [73] Muralidharan SV, Bhadury J, Nilsson LM, Green LC, McLure KG, Nilsson JA. BET bromodomain inhibitors synergize with ATR inhibitors to induce DNA damage, apoptosis, senescence-associated secretory pathway and ER stress in Myc-induced lymphoma cells. *Oncogene* 2016;35(36):4689–97.

- [74] Pires IM, Olcina MM, Anbalagan S, et al. Targeting radiation-resistant hypoxic tumour cells through ATR inhibition. *Br J Cancer* 2012;107(2):291–9.
- [75] Ma J, Li X, Su Y, et al. Mechanisms responsible for the synergistic antileukemic interactions between ATR inhibition and cytarabine in acute myeloid leukemia cells. *Sci Rep* 2017;7:41950.
- [76] Kim HJ, Min A, Im SA, et al. Anti-tumor activity of the ATR inhibitor AZD6738 in HER2 positive breast cancer cells. *Int J Cancer* 2017;140(1):109–19.
- [77] Morgado-Palacin I, Day A, Murga M, et al. Targeting the kinase activities of ATR and ATM exhibits antitumoral activity in mouse models of MLL-rearranged AML. *Sci Signal* 2016;9(445):ra91.
- [78] Zhang Y, Lai J, Du Z, et al. Targeting radioresistant breast cancer cells by single agent CHK1 inhibitor via enhancing replication stress. *Oncotarget* 2016;7(23):34688–702.
- [79] Magnussen GI, Emilsen E, Giller Fleten K, et al. Combined inhibition of the cell cycle related proteins Wee1 and Chk1/2 induces synergistic anti-cancer effect in melanoma. *BMC Cancer* 2015;15:462.
- [80] Yuan LL, Green A, David L, et al. Targeting CHK1 inhibits cell proliferation in FLT3-ITD positive acute myeloid leukemia. *Leuk Res* 2014;38(11):1342–9.
- [81] Kim H, George E, Ragland RL, et al. Targeting the ATR/CHK1 axis with PARP inhibition results in tumor regression in BRCA mutant ovarian cancer models. *Clin Cancer Res* 2016;23(12):3097–108.
- [82] Sakurikar N, Thompson R, Montano R, Eastman A. A subset of cancer cell lines is acutely sensitive to the Chk1 inhibitor MK-8776 as monotherapy due to CDK2 activation in S phase. *Oncotarget* 2016;7(2):1380–94.
- [83] Zhao J, Niu X, Li X, et al. Inhibition of CHK1 enhances cell death induced by the Bcl-2-selective inhibitor ABT-199 in acute myeloid leukemia cells. *Oncotarget* 2016;7(23):34785–99.
- [84] Duan L, Perez RE, Hansen M, Gitelis S, Maki CG. Increasing cisplatin sensitivity by schedule-dependent inhibition of AKT and Chk1. *Cancer Biol Ther* 2014;15(12):1600–12.
- [85] Sarmiento LM, Pova V, Nascimento R, et al. CHK1 overexpression in T-cell acute lymphoblastic leukemia is essential for proliferation and survival by preventing excessive replication stress. *Oncogene* 2015;34(23):2978–90.
- [86] Chila R, Basana A, Lupi M, et al. Combined inhibition of Chk1 and Wee1 as a new therapeutic strategy for mantle cell lymphoma. *Oncotarget* 2015;6(5):3394–408.
- [87] Bryant C, Rawlinson R, Massey AJ. Chk1 inhibition as a novel therapeutic strategy for treating triple-negative breast and ovarian cancers. *BMC Cancer* 2014;14:570.
- [88] Zhou ZR, Yang ZZ, Wang SJ, et al. The Chk1 inhibitor MK-8776 increases the radiosensitivity of human triple-negative breast cancer by inhibiting autophagy. *Acta Pharmacol Sin* 2017;.
- [89] Hall AB, Newsome D, Wang Y, et al. Potentiation of tumor responses to DNA damaging therapy by the selective ATR inhibitor VX-970. *Oncotarget* 2014;5(14):5674–85.
- [90] Vendetti FP, Lau A, Schamus S, Conrads TP, O'Connor MJ, Bakkenist CJ. The orally active and bioavailable ATR kinase inhibitor AZD6738 potentiates the anti-tumor effects of cisplatin to resolve ATM-deficient non-small cell lung cancer *in vivo*. *Oncotarget* 2015;6(42):44289–305.
- [91] Mohni KN, Thompson PS, Luzwick JW, et al. A synthetic lethal screen identifies DNA repair pathways that sensitize cancer cells to combined ATR inhibition and cisplatin treatments. *PLoS ONE* 2015;10(5):e0125482.
- [92] Josse R, Martin SE, Guha R, et al. ATR inhibitors VE-821 and VX-970 sensitize cancer cells to topoisomerase I inhibitors by disabling DNA replication initiation and fork elongation responses. *Cancer Res* 2014;74(23):6968–79.
- [93] Chaudhuri L, Vincelette ND, Koh BD, et al. CHK1 and WEE1 inhibition combine synergistically to enhance therapeutic efficacy in acute myeloid leukemia *ex vivo*. *Haematologica* 2014;99(4):688–96.
- [94] Russell MR, Levin K, Rader J, et al. Combination therapy targeting the Chk1 and Wee1 kinases shows therapeutic efficacy in neuroblastoma. *Cancer Res* 2013;73(2):776–84.
- [95] Mak JP, Man WY, Chow JP, Ma HT, Poon RY. Pharmacological inactivation of CHK1 and WEE1 induces mitotic catastrophe in nasopharyngeal carcinoma cells. *Oncotarget* 2015;6(25):21074–84.
- [96] Dietlein F, Kalb B, Jokic M, et al. A synergistic interaction between Chk1- and MK2 inhibitors in KRAS-mutant cancer. *Cell* 2015;162(1):146–59.
- [97] Sausville E, Lorusso P, Carducci M, et al. Phase I dose-escalation study of AZD7762, a checkpoint kinase inhibitor, in combination with gemcitabine in US patients with advanced solid tumors. *Cancer Chemother Pharmacol* 2014;73(3):539–49.
- [98] Seto T, Esaki T, Hirai F, et al. Phase I dose-escalation study of AZD7762 alone and in combination with gemcitabine in Japanese patients with advanced solid tumours. *Cancer Chemother Pharmacol* 2013;72(3):619–27.

- [99] Doi T, Yoshino T, Shitara K, et al. Phase I study of LY2603618, a CHK1 inhibitor, in combination with gemcitabine in Japanese patients with solid tumors. *Anticancer Drugs* 2015;26(10):1043–53.
- [100] Hong D, Infante J, Janku F, et al. Phase I study of LY2606368, a checkpoint kinase 1 inhibitor, in patients with advanced cancer. *J Clin Oncol* 2016;34(15):1764–71.
- [101] Karp JE, Thomas BM, Greer JM, et al. Phase I and pharmacologic trial of cytosine arabinoside with the selective checkpoint 1 inhibitor Sch 900776 in refractory acute leukemias. *Clin Cancer Res* 2012;18(24):6723–31.
- [102] Daud AI, Ashworth MT, Strosberg J, et al. Phase I dose-escalation trial of checkpoint kinase 1 inhibitor MK-8776 as monotherapy and in combination with gemcitabine in patients with advanced solid tumors. *J Clin Oncol* 2015;33(9):1060–6.
- [103] Scagliotti G, Kang JH, Smith D, et al. Phase II evaluation of LY2603618, a first-generation CHK1 inhibitor, in combination with pemetrexed in patients with advanced or metastatic non-small cell lung cancer. *Invest New Drugs* 2016;34(5):625–35.
- [104] Laquente B, Lopez-Martin J, Richards D, et al. A phase II study to evaluate LY2603618 in combination with gemcitabine in pancreatic cancer patients. *BMC Cancer* 2017;17(1):137.
- [105] Al-Ahmadie H, Iyer G, Hohl M, et al. Synthetic lethality in ATM-deficient RAD50-mutant tumors underlies outlier response to cancer therapy. *Cancer Discov* 2014;4(9):1014–21.
- [106] Hickson I, Zhao Y, Richardson CJ, et al. Identification and characterization of a novel and specific inhibitor of the ataxia-telangiectasia mutated kinase ATM. *Cancer Res* 2004;64(24):9152–9.
- [107] Golding SE, Rosenberg E, Valerie N, et al. Improved ATM kinase inhibitor KU-60019 radiosensitizes glioma cells, compromises insulin, AKT and ERK prosurvival signaling, and inhibits migration and invasion. *Mol Cancer Ther* 2009;8(10):2894–902.
- [108] Biddlestone-Thorpe L, Sajjad M, Rosenberg E, et al. ATM kinase inhibition preferentially sensitizes p53-mutant glioma to ionizing radiation. *Clin Cancer Res* 2013;19(12):3189–200.
- [109] Batey MA, Zhao Y, Kyle S, et al. Preclinical evaluation of a novel ATM inhibitor, KU59403, in vitro and in vivo in p53 functional and dysfunctional models of human cancer. *Mol Cancer Ther* 2013;12(6):959–67.
- [110] Blake SM, Stricker SH, Halavach H, et al. Inactivation of the ATMIN/ATM pathway protects against glioblastoma formation. *Elife* 2016;5. e08711.
- [111] Lin YF, Shih HY, Shang Z, Matsunaga S, Chen BP. DNA-PKcs is required to maintain stability of Chk1 and Claspin for optimal replication stress response. *Nucleic Acids Res* 2014;42(7):4463–73.
- [112] Buisson R, Boisvert JL, Benes CH, Zou L. Distinct but concerted roles of ATR, DNA-PK, and Chk1 in counteracting replication stress during S phase. *Mol Cell* 2015;59(6):1011–24.
- [113] Sanjiv K, Hagenkort A, Calderon-Montano JM, et al. Cancer-specific synthetic lethality between ATR and CHK1 kinase activities. *Cell Rep* 2016;17(12):3407–16.
- [114] Gilad O, Nabet BY, Ragland RL, et al. Combining ATR suppression with oncogenic Ras synergistically increases genomic instability, causing synthetic lethality or tumorigenesis in a dosage-dependent manner. *Cancer Res* 2010;70(23):9693–702.
- [115] Teng PN, Bateman NW, Darcy KM, et al. Pharmacologic inhibition of ATR and ATM offers clinically important distinctions to enhancing platinum or radiation response in ovarian, endometrial, and cervical cancer cells. *Gynecol Oncol* 2015;136(3):554–61.
- [116] Leszczynska KB, Dobrynin G, Leslie RE, et al. Preclinical testing of an Atr inhibitor demonstrates improved response to standard therapies for esophageal cancer. *Radiother Oncol* 2016;121(2):232–8.
- [117] Menezes DL, Holt J, Tang Y, et al. A synthetic lethal screen reveals enhanced sensitivity to ATR inhibitor treatment in mantle cell lymphoma with ATM loss-of-function. *Mol Cancer Res* 2015;13(1):120–9.
- [118] Huntoon CJ, Flatten KS, Wahner Hendrickson AE, et al. ATR inhibition broadly sensitizes ovarian cancer cells to chemotherapy independent of BRCA status. *Cancer Res* 2013;73(12):3683–91.
- [119] Prevo R, Fokas E, Reaper PM, et al. The novel ATR inhibitor VE-821 increases sensitivity of pancreatic cancer cells to radiation and chemotherapy. *Cancer Biol Ther* 2012;13(11):1072–81.
- [120] Fokas E, Prevo R, Pollard JR, et al. Targeting ATR *in vivo* using the novel inhibitor VE-822 results in selective sensitization of pancreatic tumors to radiation. *Cell Death Dis* 2012;3:e441.
- [121] Wang WT, Catto JW, Meuth M. Differential response of normal and malignant urothelial cells to CHK1 and ATM inhibitors. *Oncogene* 2015;34(22):2887–96.
- [122] Tang Y, Hamed HA, Poklepovic A, Dai Y, Grant S, Dent P. Poly(ADP-ribose) polymerase 1 modulates the lethality of CHK1 inhibitors in mammary tumors. *Mol Pharmacol* 2012;82(2):322–32.

- [123] Martino-Echarri E, Henderson BR, Brocardo MG. Targeting the DNA replication checkpoint by pharmacologic inhibition of Chk1 kinase: a strategy to sensitize APC mutant colon cancer cells to 5-fluorouracil chemotherapy. *Oncotarget* 2014;5(20):9889–900.
- [124] Origanti S, Cai SR, Munir AZ, White LS, Piwnica-Worms H. Synthetic lethality of Chk1 inhibition combined with p53 and/or p21 loss during a DNA damage response in normal and tumor cells. *Oncogene* 2013;32(5):577–88.
- [125] Gadhikar MA, Sciuto MR, Alves MV, et al. Chk1/2 inhibition overcomes the cisplatin resistance of head and neck cancer cells secondary to the loss of functional p53. *Mol Cancer Ther* 2013;12(9):1860–73.
- [126] Barker HE, Patel R, McLaughlin M, et al. CHK1 inhibition radiosensitizes head and neck cancers to paclitaxel-based chemoradiotherapy. *Mol Cancer Ther* 2016;15(9):2042–54.
- [127] Bartucci M, Svensson S, Romania P, et al. Therapeutic targeting of Chk1 in NSCLC stem cells during chemotherapy. *Cell Death Differ* 2012;19(5):768–78.
- [128] Zemanova J, Hylse O, Collakova J, et al. Chk1 inhibition significantly potentiates activity of nucleoside analogs in TP53-mutated B-lymphoid cells. *Oncotarget* 2016;7(38):62091–106.
- [129] Pei XY, Dai Y, Felthousen J, et al. Circumvention of Mcl-1-dependent drug resistance by simultaneous Chk1 and MEK1/2 inhibition in human multiple myeloma cells. *PLoS ONE* 2014;9(3):e89064.
- [130] Dai Y, Chen S, Shah R, et al. Disruption of Src function potentiates Chk1-inhibitor-induced apoptosis in human multiple myeloma cells *in vitro* and *in vivo*. *Blood* 2011;117(6):1947–57.
- [131] Cole KA, Huggins J, Laquaglia M, et al. RNAi screen of the protein kinome identifies checkpoint kinase 1 (CHK1) as a therapeutic target in neuroblastoma. *Proc Natl Acad Sci USA* 2011;108(8):3336–41.
- [132] Sankunny M, Parikh RA, Lewis DW, Gooding WE, Saunders WS, Gollin SM. Targeted inhibition of ATR or CHEK1 reverses radioresistance in oral squamous cell carcinoma cells with distal chromosome arm 11q loss. *Genes Chromosomes Cancer* 2014;53(2):129–43.
- [133] Vance S, Liu E, Zhao L, et al. Selective radiosensitization of p53 mutant pancreatic cancer cells by combined inhibition of Chk1 and PARP1. *Cell Cycle* 2011;10(24):4321–9.
- [134] Engelke CG, Parsels LA, Qian Y, et al. Sensitization of pancreatic cancer to chemoradiation by the Chk1 inhibitor MK8776. *Clin Cancer Res* 2013;19(16):4412–21.
- [135] McNeely S, Conti C, Sheikh T, et al. Chk1 inhibition after replicative stress activates a double strand break response mediated by ATM and DNA-dependent protein kinase. *Cell Cycle* 2010;9(5):995–1004.
- [136] Calvo E, Braiteh F, Von Hoff D, et al. Phase I study of CHK1 inhibitor LY2603618 in combination with gemcitabine in patients with solid tumors. *Oncology* 2016;91(5):251–60.